

# The Pediatrician's Role in Encouraging Exclusive Breastfeeding

Maya Bunik, MD, MSPH\*

\*Department of Pediatrics, University of Colorado School of Medicine, Children's Hospital Colorado, Aurora, CO

## Practice Gap

Although most pregnant women in the United States plan to breastfeed, there is a clear gap between the proportion of women who prenatally intend to breastfeed and those who actually meet their goals postpartum. (1,2) Pediatricians are often challenged by providing support and management for nursing issues.

## Objectives After completing this article, readers should be able to:

1. Describe differences between human milk and formula and the practice of breastfeeding and formula feeding.
2. Identify the factors that interfere with breastfeeding.
3. Discuss management of maternal nipple pain, low milk supply, late preterm infants, infants with allergic colitis, and problems with maternal milk oversupply.
4. Learn the best ways to support the mother-baby dyad through the birth, hospital, and postpartum periods.
5. Recognize the effects of maternal ingestion of drugs and other substances.

## INTRODUCTION

According to the US Centers for Disease Control and Prevention (CDC), many mothers initiate breastfeeding at birth, but only 22% of infants are exclusively breastfed for 6 months, and only 29% experience any breastfeeding in the first 12 months. (3) This falls short of the Healthy People 2020 goals of 25.5% exclusively breastfeeding for 6 months and 34.1% breastfeeding in the first 12 months (Fig 1). (4) Disparities exist, including differences related to race and ethnicity. For example, Hispanics have higher breastfeeding initiation and duration rates than the black population but lower exclusivity rates than the white population. (5) Moreover, lower breastfeeding rates are also associated with socioeconomic disparities, such as lower education level, poverty, young age, high body mass index, not abstaining from alcohol use, smoking within 6 weeks of delivery, and living in rural areas. These disparities in breastfeeding affect subsequent child health.

**AUTHOR DISCLOSURE** Dr Bunik has disclosed no financial relationships relevant to this article. This commentary does not contain a discussion of an unapproved/investigative use of a commercial product/device.

### ABBREVIATIONS

BFHI Baby-Friendly Hospital Initiative  
CDC Centers for Disease Control and Prevention  
THC tetrahydrocannabinol

The US Preventive Services Task Force recently gave primary care breastfeeding support a “B” recommendation, meaning there is strong evidence that office-based health care promotion should be a priority. (6) Therefore, pediatricians, as advocates and partners for the health of children, should support and promote breastfeeding in mother-infant dyads. Most current intervention efforts are not adequately communicating to families the importance of breastfeeding exclusivity and duration as the norm for all mother-infant dyads. (7,8) Professional education in breastfeeding and helping mothers get access to professional support are provider responsibilities highlighted in the CDC’s “Guide to Strategies to Support Breastfeeding Mothers and Babies.” (9) In addition, providers should make sure that support for breastfeeding starts in the hospital at birth and then continues with office-based support and peer support at discharge. If a mother is returning to work, she will need support from her employer, as well as her childcare provider.

In this review article, the main objective is to present ways for pediatricians to address the current continued barriers for exclusive breastfeeding, to describe early management challenges, and to provide key tools to foster the rewarding relationship that results when pediatricians and/or staff in their practice play a main role in breastfeeding success.

## UNDERSTANDING THE DIFFERENCES BETWEEN HUMAN MILK AND FORMULA

### Content Comparison

Human milk is superior to formula because of the immunologic properties that are passed on to the infant and are

the basis for disease prevention. Human milk has bioactive factors such as living cells, enzymes, and antibodies that offer immune protection and support the physiological microbes of the gastrointestinal tract. Tow stated in her breastfeeding review that the “most significant inheritance a child will ever receive is the maternal microbiome.” (10) Immunoglobulins in human milk are predominantly secretory immunoglobulin A, with smaller amounts of immunoglobulin M and immunoglobulin G. Cells in the breasts interact with maternal plasma from the bronchial tree and intestine to produce immunoglobulin A antibodies that offer specific protection against pathogens in the mother’s environment.

High concentration and structural diversity of human milk oligosaccharides, as a group of more than 200 identified complex and diverse glycans, are resistant to gastrointestinal digestion and reach the colon as the first prebiotics. Many human milk oligosaccharides are known to directly interact with the surface of pathogenic bacteria, such as *Haemophilus influenzae* and *Streptococcus pneumoniae*, and inhibit binding and toxins to the host receptors. (11)

In this way, human milk oligosaccharides act as decoys to protect infants from infectious diseases. They are also responsible for a diverse spectrum of functions that include the compositional development of gut microbiota, prevention of intestinal infections, and development of the brain. (12) Lactoferrin, lysozyme, complement,  $\alpha$ -lactalbumin, and casein are other important bioactive proteins that act in concert with the complex immune framework. (13)

The composition of human milk varies somewhat from feeding to feeding but usually ranges from 19 to 21 calories per ounce. Volumes can vary throughout the day and also from mother to mother.




Figure 1. CDC graph of the percentage of babies breastfeeding during the first year.

Formula companies are constantly improving on nutritional content to make formula closer to human milk—for example, by adding docosahexaenoic acid and probiotics. Unfortunately, various labeling and advertising on their products can lead families to believe that certain formulas can eliminate infants' gastrointestinal symptoms, such as gas and reflux. In terms of other nutritional components, formulas contain slightly higher levels of protein than human milk (1.5 g vs 1.1 g per 110 mL, respectively).

### Infant Early Weight Loss Nomograms

In a large Kaiser study, investigators found that 5% of exclusively breastfed, vaginally delivered newborns and 10% of cesarean-delivered newborns lost 10% or more of their birth weight 48 hours after delivery. Formula-fed newborns had much lower weight loss rates of 2.9% and 3.5%, respectively. These nomograms (see Fig 2) should be used for early identification of newborns on a trajectory of greater weight loss and follow-up for associated morbidities. (14,15)

### Return to Birth Weight and Weight Gain

Some exclusively breastfed newborns may require slightly longer than 2 weeks to return to their birth weight. (16) Close monitoring with frequent weight checks is preferable in this situation because the addition of supplementation may cause more morbidity than watchful waiting.

### Infant Temperament and Maternal Bonding

Mothers often report that breastfeeding is an enjoyable bonding time with their infants. (17) Functional brain magnetic resonance imaging has been used to compare exclusively breastfeeding mothers with exclusively formula-feeding mothers as they listened to their infants' own cries versus a control infant's cry. Breastfeeding mothers showed greater brain activations on images while listening to their own baby cry. Studies have also supported the fact that breastfeeding mothers have a higher-rated sensitivity score and are more in tune with infant temperament than mothers who are formula feeding. (18,19)

### Milk Volumes

Volume intake varies somewhat with breastfed infants because of the diurnal nature of maternal milk supply (with higher prolactin levels at night). In formula-fed infants, the bottle feedings are controlled by the parent or caregiver, who determines volume, frequency, hunger, and satiety cues. Conversely, infants who are actively feeding and are not sleepy at the breast are in control of the volume of the milk that they transfer from the mother.

### Stool Production Patterns

Stool production patterns depend on adequate human milk intake and age of a baby. Expect one stool for each day of life until day 4, when a mother's milk is fully in, leading to a transition in stool color, from meconium black to green to yellow during this time. In the first few weeks, stools commonly occur with every feeding or every other feeding. The "breastfed stool variant" occurs in about one-third of breastfed infants starting at 4 to 6 weeks of age. (20) One soft, voluminous stool occurs usually every 3 to 4 days (but can take up to 12–14 days), and this is thought to be due to almost complete absorption of human milk. At times, the infant can appear uncomfortable, but more commonly, the infant is asymptomatic. It is not known why this occurs in only some breastfed infants.

### Vitamin D Supplementation

Vitamin D is generated through the skin from exposure to the sun, and owing to concurrent recommendations for judicious use of sunscreen and sun avoidance, breastfeeding infants require 400 IU of vitamin D supplementation per day. However, adherence to these recommendations has been low. (21,22) Recent work suggests that maternal vitamin D at levels of 6400 IU per day are adequate for transfer of vitamin D to the infant. (23) Most infant formulas contain the added 400 IU of vitamin D with an intake range of 26 to 32 oz.

### Developmental Phases

Developmental phases can sometimes interfere with breastfeeding and may differ from infants who are bottle-fed human milk or formula fed. At 4 to 5 months of age, when the infant's vision improves, most infants can become distracted during a feeding, and it can be challenging to keep the infant on the task of completing a full feeding. This may result in more hunger, more frequent feedings, more feeding in the nighttime, and, in rare cases, slower weight gain. Nursing in a quiet, dark room with minimal distractions can help. When infants begin teething, biting can also become a problem. Mechanically, the infant cannot bite and ingest milk at the same time. Infants who bite at the breast when the breast is offered are usually not ready to feed, and biting at the end of the nursing session may mean that they are full. Anticipating this biting behavior may be helpful, and disengaging the infant quickly and ending the feeding usually sends the message that this is an undesirable behavior. Again, perception that the infant is not satisfied by breast milk alone is cited consistently as one of the top reasons for stopping breastfeeding, regardless of weaning





Figure 2. Nomograms for early neonate weight loss (14). A. Estimated percentile curves of percentage weight loss according to time after birth for vaginal deliveries. B. Estimated percentile curves of percentage weight loss according to time after birth for cesarean deliveries.

age. (24) As infants become older, they are usually more effective at breastfeeding and can transfer milk in less than 5 minutes. Mothers' breasts feel emptier, too, but the infant can usually extract the milk even if the suckling seems less effective. This concern about decrease in milk supply is a common misconception with older infants at the 9-month milestone. Providing reassurance at all of these phases can be comforting to the mother and prevent early cessation of breastfeeding.

### Risk of not Breastfeeding for Infant and Mother

The dose-responsive benefits of breastfeeding for both the infant and mother are well known and well established and should be the basis for encouraging mothers to breastfeed exclusively. (25,26) The Surgeon General's Report in 2011 and others suggest reframing this argument as the "risk of not breastfeeding." (27,28) The risks of not breastfeeding for infants are shown in Table 1. Mothers who do not breastfeed do not return to prepregnancy weight as quickly and do not have the associated decreased incidence of type 2 diabetes, osteoporosis, hypertension, heart attack, stroke, postpartum depression, breast cancer, and ovarian cancer, as do mothers who breastfeed. (29-34)

## ADDRESSING BARRIERS TO EXCLUSIVE BREASTFEEDING

### Pain and Low Milk Supply

Pain, sore nipples, and perceived or real insufficient milk supply are the main reasons for early cessation of breastfeeding. (24,35) Sore nipples occur in about one-third of mothers. (36) To avoid pain and trauma, it is best to ensure proper latching by encouraging the newborn to open the mouth wide by tickling his or her lips with a finger or nipple. The mother should pull the baby in close to the mother's abdomen and support the back, so that the newborn's chin dives into the breast and the newborn's nose touches the breast at the nipple. Once latched, the baby's lips should be untucked and flared, like "fish lips." (Fig 3). Although neonates may spend up to 45 to 60 minutes nursing at a time, some of this time may not be nutritive, which is referred to as sleepy "flutter feeding." Prolonged time spent suckling can result in nipple soreness in the early days of breastfeeding, so limiting the time to 30 minutes total, while keeping the neonate on task by tickling or keeping his or her arm up in the air, helps. (20) The mother should break the newborn's strong mouth suction by putting her finger in the corner of the baby's mouth. If the mother has nipple redness or cracks, she should find some relief by applying

TABLE 1. **Highlighted Risks of Not Breastfeeding for Infants**

OUTCOME	EXCESS RISK, %
Full-term infants	
Acute ear infection (otitis media)	100
Eczema (atopic dermatitis)	47
Diarrhea and vomiting (gastrointestinal infection)	178
Hospitalization for lower respiratory tract diseases in the first year	257
Asthma, with family history	67
Asthma, no family history	35
Childhood obesity	32
Type 2 diabetes mellitus	64
Acute lymphocytic leukemia	23
Acute myelogenous leukemia	18
Sudden infant death syndrome	56
Preterm infants	
Necrotizing enterocolitis	138

Adapted from US Department of Health & Human Services. The Surgeon General's Call to Action to Support Breastfeeding. Accessed at <http://www.surgeongeneral.gov/library/calls/breastfeeding/> on June 15, 2016 (27)

lanolin, as well as hydrogel or soothing pads. Most postpartum nipple discomfort usually improves by day 7 to day 10. A quick list of common causes of sore nipples is shown in Table 2. (20)

Many mothers perceive that human milk coming from the breasts is not enough to exclusively feed or satisfy an infant. Pediatricians should address this concern early on. At first, human milk comes in small yellow volumes as colostrum, and then in 3 to 4 days, the milk becomes more white and watery in appearance. Most women have a hard time believing that all the nutrition a baby needs can come from her breast, and formula is a readily available option when there is maternal doubt. (37,38) Causes for irreversible low milk production include primary glandular insufficiency (<5% of women), previous breast surgeries and associated scarring, and severe postpartum birth complications, which usually involve hypertension or blood loss. Evaluation for poor latch, sleepy behavior at the breast, and inadequate milk removal will help identify the most likely reversible causes for milk supply concerns. Occasionally, oral contraceptives or pseudoephedrine decongestants

can cause a decrease in milk supply. Frequent weight checks in the office, pre- and postfeeding weight checks on a sensitive scale (a scale made to weigh to the nearest gram), and an early-morning pumping session are ways to assess and reassure mothers about milk supply. It is critical to address all of these milk supply issues so as to not unnecessarily disrupt the path toward exclusive breastfeeding.

### Breastfeeding Intention and Self-efficacy

Studies continue to show that mothers who have a strong intention to breastfeed prenatally are more likely to achieve their breastfeeding goals. They are more likely to overcome some variables that could affect their success, such as pain, fear of difficulty, birth method, partner support, and even medical complications. However, according to the latest infant feeding study, two-thirds of mothers who intend to exclusively breastfeed are not meeting their intended duration. (39) Increased Baby-Friendly Hospital Initiative (BFHI) practices, particularly giving neonates only breast milk in the hospital, may help more mothers achieve their exclusive breastfeeding intentions. (39) Prenatal maternal knowledge about infant health benefits and developing a level of comfort with breastfeeding in social settings was found to be directly related to the intention to exclusively breastfeed. (40,41) Numerous studies by Dennis and

colleagues in the past decade have shown that having early breastfeeding confidence in the first weeks is associated with longer breastfeeding duration. The validated short form, 14-item Breastfeeding Self-Efficacy Scale (42), includes statements such as, “I can always successfully cope with breastfeeding like I have other challenging tasks,” and can be used to evaluate maternal level of confidence with breastfeeding.

## IMPROVING ACCESS TO PROFESSIONAL SUPPORT

### Birth Hospital Practices Matter

Hospital practices supportive of breastfeeding in the hours and days after birth make a difference in breastfeeding rates. Currently, 18% of US births occur in 11% of BFHI-designated facilities. (43) Yotebieng et al (44) showed that the number of specific BFHI practices had a cumulative effect on exclusivity of breastfeeding. To achieve breastfeeding exclusivity, hospitals should be encouraged to adopt some of these BFHI practices, if not all of them.

The following are the BFHI Ten Steps associated with increased breastfeeding exclusivity:

**Written breastfeeding policy.** The process of developing policy brings clarification to all staff levels. One example of this success is the decline in distribution of infant formula company discharge bags and sample packs over the past 9 years. (45)

**Staff competency assessment.** All staff should be trained in the skills and messaging necessary to support new mothers and breastfeeding.

**Prenatal breastfeeding education.** To affect maternal knowledge and intent as described previously, providing education prior to delivery can help mothers overcome the unexpected challenges of early latch and breastfeeding. One study indicated that 17% of mothers reported that their physician, nurse, or other health care worker missed the opportunity to talk about breastfeeding during any of their prenatal visits. (46)

**Early initiation for latch, skin-to-skin kangaroo care, and nursing immediately after birth.** Newborns who nurse in the first hours after birth appear to be more successful with latching and nursing later on, and putting the newborn on the mother skin to skin cues the mother and newborn to nurse. (47) Early work on skin-to-skin contact indicates that it may lead to exclusive breastfeeding. (48)

**Teaching breastfeeding techniques.** Whenever possible, the hospital staff should review what mothers learned prenatally. Observing the mother and practicing latch and



Figure 3. A good, wide, open latch is shown with flared lips.



TABLE 2. Quick Reference for Pain with Breastfeeding (20)

INQUIRY	CAUSE OF PAIN	RECOMMENDATION
Lips tucked under—"grandpa lips"? Not opening the mouth wide enough and getting only the nipple in the mouth?	Poor latch	Untuck the lips Wait for a wide-open mouth; may need the baby to start feeding before becoming too awake and hungry to increase cooperation
Early days' discomfort from baby's vacuum suction? Any blanching?	Discomfort in the first weeks vs high suckling pressure	Lanolin  Deep breathing Review of good latch
Blister-like lesions on breast?	Herpes	Avoid nursing on the affected side
Pink-tinged nipples? Itching? Shooting pain in the breast?	<i>Candida</i> infection	Simultaneous antifungal treatment of mother and baby (all-purpose nipple ointment <sup>a</sup> not adequate)
Does baby's tongue extend beyond the gums? Does baby's tongue move up and sideways when you rub the gums?	Tongue-tie, other mouth abnormalities	If any suspicion, get a formal evaluation  (The author's experience is that tongue-tie is currently a popular overdiagnosis)
Shiny white dot on the tip of the nipple?	Bleb	Open up with a sterile needle; has a high rate of reoccurrence
Dry, flaky rash?  History of allergies or eczema?	Eczema or irritant dermatitis	Apply over-the-counter hydrocortisone and, if there's no improvement, may need a more potent version via prescription May have allergy to lanolin, detergents/bleach, soaps
Sensitivity of nipples to cold or stimulation? Color change of nipple after nursing?	Vasospasm of nipple/Raynaud phenomenon	Needs evaluation, will likely need prescription for nifedipine
Plentiful milk supply? Baby pulls off with squirts of milk a few minutes into a nursing session?	Clamping down due to oversupply	Lean back with nursing, because it affords baby better control of fast flow
Soreness beyond the nipple? Area of redness of the breast? Fever?	Mastitis	Needs evaluation, will likely need antibiotics

<sup>a</sup>All-purpose nipple ointment consists of compounded antibiotic, antifungal, and anti-inflammatory ointments.

techniques to keep the newborn on task when breastfeeding empowers the mother to continue nursing after discharge from the hospital.

**Limiting non-human milk feedings.** Modeling the hospital use of mother's own milk whenever possible sends a strong message to families, such that they are less likely to use formula supplementation when they go home from the hospital. Although some birth hospitals are providing donor milk to term neonates, (49) there are no studies to support this practice, which is costly and not covered by insurance. In high-risk neonates, donor human milk is best provided from an established not-for-profit milk bank

that is part of the Human Milk Banking Association of North America. (50) Donors are not paid and are carefully screened. Milk is collected according to guidelines and pasteurized. Although pasteurization provided by milk banks may affect some of the immune properties of human milk, much is still preserved. Milk sharing or purchasing on the Internet should be avoided, since there is potential for infections or contamination of milk with drugs. (51-54)

**Rooming-in, including performing milk expression with the newborn nearby.** Newborns and their mothers should be kept in close proximity for as much time as possible in the

early breastfeeding days. (55) The newborn should also be close by when the mother uses a breast pump. (56) Neonate safety should be considered when there are medical issues with the maternal-infant dyad.

**Teaching feeding cues.** Getting the newborn to the breast early, usually every 1.5 to 2 hours, can help the latch process be less stressful for both mother and baby. Crying is a late hunger cue that can add to everyone's frustrations and desire to stop the crying with bottle feeding, thereby giving the newborn unnecessary supplementation.

**Limiting pacifier use.** The American Academy of Pediatrics suggests that there are exceptions to limiting pacifier use, such as when an infant has to undergo painful procedures and when an infant needs calming. It is best to introduce pacifiers after breastfeeding is established, so that feedings are not missed in the early weeks. (57)

**Postdischarge support.** After hospital discharge, mothers need to know who to call if they have questions, as well as have a plan for routine follow-up checks and breastfeeding problems. Postdischarge support is an important "outpatient extension" of BFHI.

### Peer-to-Peer Support

Peer support by "mentor mothers" who have breastfed and are from the same community or ethnic group can be provided in several ways, including support groups (eg, La Leche League International or the birth hospital) and one-on-one support through telephone calls or visits in a home or clinical setting. Systematic reviews have demonstrated peer counselor programs to be effective in increasing breastfeeding exclusivity. (58,59) Women who provide peer support receive specific training for the best ways to offer emotional support, encouragement, education, and help with breastfeeding problems. (60,61)

### Breastfeeding-Friendly Office Practice

The steps for creating a breastfeeding-friendly office practice are shown in Fig 4.

**Evaluate early and often in the first weeks.** Maternal milk supply is established in the first 3 to 4 weeks postpartum, thereby mandating close evaluation and follow-up to ensure continued breastfeeding. Screening questions (Fig 5) were developed for telephone triage and can be used to determine the level of concern for urgent evaluation. Breastfeeding difficulties usually cause substantial distress in a new family, so determining the level of urgency can help with scheduling an earlier assessment if needed. If there is uncertainty about milk supply or problems with latch, the mother

should be seen in the pediatric office or by an outside lactation specialist.

**Decide about supplementation.** Supplementation with human milk is preferable to formula in times when there is concern about infant weight gain, because even a small amount of cow milk can disrupt the intestinal microbe environment. (62) Moreover, some infants can develop a preference early on in the feeding process, since the bottle method of feeding has a faster flow than the breast. "Slow-flow" nipples (which do not drip when the bottle is turned upside down) can help, but feeding from the breast is commonly slower than any bottle system.

**Address combination feeding.** In some cultures, the issues of combination feeding are widely accepted and associated with acculturation in the United States. "Las dos cosas" (literally, "those two things," or both breast milk and bottle formula) has been a well-described practice, particularly in Latinas, because mothers want their infants to have the "best of both worlds." (63) In addition, there is a common perception that breast milk is not enough for the infant after 3 months of age. (64,65) Meta-analyses have demonstrated that interventions to encourage exclusive breastfeeding among Latinas should ideally begin in the prenatal setting and involve frequent contact, especially with an International Board-Certified Lactation Consultant. (66)

**Advocate for maternity leave and support the return-to-work transition.** Studies show that women who intend to return to work within a year of their child's birth are less likely to initiate breastfeeding, and those who work full-time tend to breastfeed for shorter periods than those who work part-time or do not work out of the home. (67,68) Not surprisingly, women in salaried jobs and those with longer maternity leaves are more likely to breastfeed. (69) Unfortunately, planning for return to work seems to affect mothers' decision-making early in the postpartum period. (70) Many women worry about having enough milk stored and having to express milk at work. A double-sided pump is the best choice for a working mother because of its efficiency. Under the Affordable Care Act of 2014, insurance companies provide pumps as a covered benefit, but anecdotal experience suggests that these pumps may not always be of the best caliber. (71) Mothers should be encouraged to talk to their employers early during pregnancy and try to take at least 6 postpartum weeks off of work, if possible. Maternity leave of up to 6 weeks compared to 6 to 12 weeks after delivery was associated with higher odds of failure to establish breastfeeding. (72) Under the Affordable Care Act, working mothers should be provided with accommodations,



## OFFERING BREASTFEEDING SUPPORT IN YOUR OFFICE IS AS EASY AS 1-2-3

#1 Train internal staff or provide early referral for breastfeeding management and support. This is also known as extending the Baby-Friendly Hospital Initiative to your office.

#2 Provide resources for mothers, such as hospital-based drop-in clinics or groups or other mother support groups. It is always a good idea to attend or have one of your staff attend these sites so that you are sure that the advice and any concerns raised are addressed appropriately and in line with general American Academy of Pediatrics recommendations.

#3 Avoid storing and giving out formula samples in your office. It may seem supportive, but it gives the wrong message about breastfeeding exclusivity. Providing information about pumps, pump rental stations, or hand expression is a better idea.

Figure 4. Offering breastfeeding support in your office is as easy as “1-2-3.”

such as time and a private 4-foot by 6-foot space in which to pump. (73)

### OTHER BREASTFEEDING CHALLENGES THAT PROVIDERS NEED TO ADDRESS

#### Oversupply

Most newborns, even when they are older, only ingest a maximum of 3.0 to 3.5 oz of milk at the breast, (74) so mothers who are producing more than that amount usually

have a state of oversupply. Not only does oversupply cause uncomfortable fullness and leakage for the mother, but milk could be continuously “leftover” in the breast, leading to possible stasis, plugged ducts, and even mastitis. Mothers with milk oversupply should lean back as much as possible when nursing so that the infant has some ability to respond better to the fast flow, which occurs with let-down. In this position, the infant is less likely to pull off of the breast. The process of slowly and carefully down-regulating the milk supply, with less pumping and trending toward one-sided

- 1. What is your baby’s age and gestational age? Was your baby born early or on time? If early, how early? Do you have discharge sheets at home? Did the hospital tell you your newborn’s gestational age?** (Late preterm is high risk.)
- 2. Is your baby acting sick or abnormal in any way (eg, weak, decreased activity)?** Rule out sepsis, particularly in babies 4 weeks of age and younger.
- 3. Is breastfeeding going well?** If not perceived as going well, mother and baby may need to be seen in the office today or tomorrow (within 24 hours).
- 4. How many times have you breastfed in the past day?** Ten to 12 times per 24 hours is optimal; 8 times is minimal. Suboptimal nursing sessions require evaluation.
- 5. How long is your baby awake and actively suckling and swallowing at the breast during a feeding?** Baby should be actively feeding at the breast without long pauses or flutter feeding for at least 10 minutes. Just as with microwave popcorn, pauses or extended time (ie, “cooking the popcorn” too long) are not effective (analogy courtesy of Sheela Geraghty, MD, MS, IBCLC, FAAP). The mother may need assistance with latch and keeping baby on task for nursing sessions.
- 6. What color are your baby’s stools?** By day 4, stools should be yellow and seedy, not black or green transitional stools.
- 7. How many stools has your baby had in the past day?** The goal is 1 stool per day of life up to day 4 (ie, by day 4, the baby should have at least 4 stools daily). A suboptimal stool pattern requires evaluation.
- 8. How many wet urine diapers has your baby had in the past day?** Seven to 8 wet urine diapers is normal (exception: 3 wet diapers per day can be normal for the first 5 days). A suboptimal urine pattern requires evaluation.
- 9. Do your breasts feel full before feedings and softer afterward?** The optimal answer is yes. Before the milk is in, most mothers will not notice any change. If baby is close to 2 weeks of age, the mother’s breasts may be adjusting to what the baby’s needs are, and she may experience only mild symptoms of engorgement.
- 10. How many times have you supplemented with formula in the past day?** Supplementation more than once in 24 hours can affect milk supply or may indicate breastfeeding difficulties.

Figure 5. Triage assessment questions are given for the early postpartum period—the first 2 weeks. When using these advice topics, it is best to begin each call with the following 10 screening questions and then ask, “What is your main breastfeeding question or concern?” Adapted from the Screening Form for Early Follow-Up of Breastfed Newborns on the Dr. Mom Web site at <http://www.dr-mom.com>. Reproduced with permission.

**TABLE 3. Touch Points for Overcoming Obstacles to Breastfeeding Exclusivity (20)**

TIME POINT	PARENTAL CONCERN	MAIN OBSTACLE	PROVIDER ADVICE
Prenatal	"I want to breastfeed, but since I am going to work, I need to be able to formula feed, too."	Lack of information about combining breastfeeding and working Lack of information about milk expression and access to breast pumps	Strongly encourage attendance at a prenatal breastfeeding class (deserves equal time to birthing class education) Consider a longer maternity leave, if possible Prepare to simplify life during the transition to parenting
	"My husband/partner and other family members will want to help feed the baby. Won't they feel excluded if I only breastfeed?"	Family members wanting to feed the baby	Enlist father's/partner's help in supporting the nursing partner; fathers/partners can interact with their newborn by holding the baby skin to skin or taking the baby out while mom sleeps After breastfeeding is well established, others can feed the baby expressed milk by bottle
	"I want to do combination feeding, or las dos cosas."	Desire for "the best of both worlds" by combination feeding Lack of knowledge about the importance of frequent and exclusive breastfeeding during the early postpartum weeks for establishing mother's milk supply	"Puro pecho," or only mother's own milk, provides greater health benefits and helps maintain an abundant milk supply If eligible, enrollment in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) offers breastfeeding mothers a substantial food package, counseling, breast pumps, and peer counselors
Birth	"My friend says it is a good idea to ask the nurses to care for my baby at night so I can get some sleep."	Unrealistic expectations for the postbirth hospital stay Lack of prenatal education	Promote immediate skin-to-skin contact after birth to facilitate the initiation of breastfeeding within the first hour Teach the mother to interpret her newborn's feeding cues and breastfeed as often as baby wants; advocate for no routine formula use in the system of care
		Frequent interruptions and excessive visitors deplete new mothers Increased risk of formula supplements for nighttime births from 9 pm to 6 am	Advise the mother to request help in the hospital with breastfeeding to promote task mastery Encourage continuous rooming-in, where the mother can practice being with her baby in a controlled setting and learn to latch baby comfortably and effectively
		Belief that the small amount of colostrum is insufficient until the "milk comes in"	Explain the potency and adequacy of colostrum and the rapid increase in milk production from 36 to 96 h
3–5 d	"Now that we are home from the hospital, the baby seems to be feeding every hour. She or he doesn't seem satisfied with breast milk alone."	Lack of knowledge about normal frequency of feedings for breastfed newborns Newborns typically begin feeding more frequently the second night after birth, when the baby is at home Concern about whether the newborn is getting enough milk, due to the mother's inability to see what the newborn takes at the breast Sleepy newborn	Explain that 8–12 feedings in 24 h are typical and necessary to establish an abundant milk supply Provide a hand-pump or teach hand expression, so the mother can see that she has milk Explain normal newborn elimination patterns once the mother's milk comes in (3–5 voids and 3–4 stools per day by 3–5 d; onset of yellow, seedy milk stools by 4–5 d) Perform newborn test weights (before and after feeding) to reassure the mother about baby's milk intake at a feeding Teach the mother the difference between newborn "flutter sucking" or "nibbling" that results in only a trickle of milk at the breast versus "drinking" milk, with active sucking and regular swallowing Tickling under the axilla or holding a hand up can help keep baby on task at the breast; or, compressing the breast when the baby stops slow, deep sucking can deliver a spray of milk to entice him or her to start drinking again Anticipate newborn appetite spurt at about 10–14 d of age

*Continued*

TABLE 3. (Continued)

TIME POINT	PARENTAL CONCERN	MAIN OBSTACLE	PROVIDER ADVICE
	"My nipples are sore and cracked. Can I take a break and give my baby a little formula?"	Sore nipples are usually attributable to incorrect latch-on technique and are a common reason that mothers discontinue breastfeeding early or start supplements	Observe a nursing session to evaluate latch; consider referring the mother to a lactation consultant for one-on-one assistance with latch
2 wk	"My breasts do not feel very full anymore. I'm afraid my milk went away."	As postpartum breast engorgement resolves and the breasts adjust to making and releasing milk, mothers may perceive that they have insufficient milk supply	Expect the newborn to be above birth weight by 10–14 d and reassure the mother about the newborn's rate of weight gain since the 3–5-d visit Although the mother's breasts are less swollen than during postpartum engorgement, they should feel fuller before feedings and softer afterward
	"How can I know my baby is getting enough?"	The 10–14-d appetite spurt can cause the mother to doubt the adequacy of her milk supply	Consider performing test weights (before and after feeding) to reassure the mother about her newborn's intake Anticipate another appetite spurt at about 3 wk of age
1 mo	"My baby is crying a lot, and I am tired and need sleep."	Normal infant crying peaks at about 6 wk (3–5 h in 24 h) Mother may attribute infant crying to hunger or an adverse reaction to her milk	Congratulate the mother on a full month of breastfeeding! If infant has gained weight appropriately, reassure the mother about the adequacy of her milk supply Offer coping strategies for infant crying, including holding the baby skin to skin; the 5 S's <sup>a</sup> (however, swaddle with the baby's hands up near the head to help assess feeding cues); use of an infant carrier; going for a stroller or car ride; observing a period of "purple crying"
	"Nothing seems to calm her/him except the bottle."	If the infant drinks milk from a bottle that is offered, the mother may assume that her infant is not satisfied by breastfeeding	Explain that infant sucking is reflexive, and drinking from an offered bottle doesn't always mean that the baby is hungry; baby "can't scream and suck at the same time," so the bottle may appear to calm baby, just as a pacifier might If mom desires to offer a bottle, use expressed milk as the supplement Forewarn the mother about cluster feedings (late afternoon/evening) and upcoming appetite spurts, occurring at about 6 wk and 3 mo
2 mo	"My mother said that, if I give my baby rice cereal in a bottle before bedtime, he or she may sleep longer at night."	Parental sleep deprivation The mother may have already returned to work, which often increases fatigue and leads to a decrease in milk supply	Explain the lack of evidence that rice cereal or other solid foods increase infant sleep Remind the mother that adding complementary foods is a project and increases the workload for parents Reinforce the benefits of exclusive breastfeeding for maternal-infant health and mother's milk supply
	"I am going back to work and am worried that I do not have enough frozen stores of milk. Are there any herbs I can take to keep my milk supply strong?"	Lack of knowledge about the principles of milk production and unrealistic beliefs about the effectiveness of galactagogues	Enlist help from others, including support for returning to work Explain that there is no "magic pill" or special tea to increase the mother's milk supply; the key to ongoing milk production is frequent, effective milk removal (every 3–4 h) Caution the mother to avoid going long intervals without draining her breasts
4 mo	"My baby seems to only eat for a few minutes, and when I try to put her/him back to the breast, she or he refuses."	Misinterpretation of infant's efficiency in nursing causes concern about infant milk intake	Explain that infants become more efficient at breastfeeding, and by 3 mo, they may drain the breast in 4–7 min Reinforce continuing to delay the introduction of solid foods

Continued

TABLE 3. (Continued)

TIME POINT	PARENTAL CONCERN	MAIN OBSTACLE	PROVIDER ADVICE
	"My baby seems more interested in everything around him or her than in nursing at the breast."	Normal infant distractibility causes the mother to believe her infant is self-weaning	Explain that distractibility is a normal developmental behavior at this age and that short, efficient feedings are common Nurse in a quiet, darkened room
6 mo	"My baby is drooling and rubbing on her or his gums all the time. I do not think that I can continue to breastfeed because my baby might bite me."	Common myth that a mother needs to wean when her baby gets teeth to avoid being bitten while breastfeeding	Congratulate the mother on 6 mo of exclusive breastfeeding! Explain that infants cannot bite and actively breastfeed at the same time. Biting tends to occur if the breast is offered when the infant is not interested or at the end of the feeding If the infant bites, say "No biting," touch the infant's lips, set the baby down, and briefly leave the room
	"My baby has refused to breastfeed for almost a whole day now. Is she or he ready to wean?"	Misinterpretation of sudden breastfeeding refusal ("nursing strike") to mean that a baby is self-weaning	Explain that some babies may suddenly refuse the breast between 4 and 7 mo of age for no apparent reason; common causes include upper respiratory infection, ear infection, teething, regular exposure to bottle-feeding, use of a new soap/perfume, maternal stress, or a decrease in milk supply Because many babies will nurse while asleep, try offering the breast when the baby is drowsy or asleep Regularly express milk if the baby won't nurse, and feed the pumped milk until the infant resumes breastfeeding

<sup>a</sup>The 5 S's consist of swaddling, side lying, swaying, shushing, and sucking. From: Karp, Harvey. *The Happiest Baby on the Block*, 2nd Ed. New York: NY: Bantam Books; 2015.

feeding, is recommended but can be challenging because it seems counterintuitive to want to make less milk.

### Tongue-Tie

It has become increasingly common for infants to have their tongue- or lip-tie diagnosed and clipped as part of a breastfeeding consultation. (75) Despite little evidence for these procedures, tongue and lip clipping have become widely accepted. (76)

Anterior tongue-tie, which causes the tongue to have decreased lateral movement and decreased frontal movement up and over the gums, can cause pain and ineffective latch when breastfeeding. Anterior tongue-tie can be associated with ineffective nursing and decreasing milk supply. (77) With posterior tongue-tie, the tongue is tacked down at the back of the mouth—a condition that is difficult to diagnose. This anatomic condition is not universally accepted by otolaryngologists and is not easily corrected with surgery, which involves releasing tissue at the base of the tongue. (78) Any infant who has undergone the procedure, by means of either incision or laser (now performed by many dentists and others), should be followed up closely. If breastfeeding does not improve after the procedure, more feeding evaluation and assistance may be needed.

### Allergic Colitis

Allergic colitis is a cell-mediated hypersensitivity disorder that affects the large intestinal tract in less than 1% of

exclusively breastfed infants. The most common symptom associated with food allergy in the infant is bloody stools. (79) Dietary proteins in the mother's milk are responsible, which are usually due to maternal ingestion of dairy products. Breastfed infants with allergic colitis are typically well appearing and rarely have reflux symptoms, such as vomiting, diarrhea, and abdominal distention. Elimination of cow milk from the maternal diet is the first step in treatment, and in most cases, symptoms in the infant should improve within 3 to 4 days. If continued mucus and blood are detected, then mothers need to eliminate soy products next, followed by the remaining causal agents: eggs, nuts, wheat, corn, strawberries, citrus, and chocolate. Compliance with these elimination diets is challenging initially and even more difficult to maintain, so mothers should be cautioned to eliminate only one item at a time to be sure that eliminating one particular food from the diet is necessary. Interestingly, most infants with allergic colitis tolerate cow milk after their first birthday. (80)

### Fussiness

Just as breastfeeding is established at the end of the first month of life, many infants manifest common fussy behaviors that may be misinterpreted as gastroesophageal reflux or food allergies. Discussion about the normal crying curve that peaks at 6 weeks and about ways to soothe a crying

infant should be part of anticipatory guidance for the breastfed infant. (81)(82) Mothers and family members (and medical providers) may erroneously blame breastfeeding in some way (diet, milk supply, medications) as a reason for the fussiness.

### Reflux

Reflux symptoms, such as increased spitting up and fussiness, can cause unnecessary supplementation in the infant. Reflux can cause abdominal pain when the infant is not upright during nursing, which could mistakenly lead to elimination diets in the mother. About 50% of infants experience effortless spitting up of small amounts of breast milk (1–2 teaspoons). This spitting up should be distinguished from vomiting, which is forcefully throwing up more volume. It can be helpful to instruct the family that the stomach is like an “untied water balloon” and that any abrupt movement can cause “leakage.” Providing the mother with reassurance that reflux will improve with age in most infants can help prevent the mother and infant from taking unnecessary paths toward medication use.

### Maternal Medications

The need to prescribe almost any type of maternal medication can lead to erroneous early cessation of breastfeeding. The decision-making process around medication use while breastfeeding should be one of joint negotiation between mother and provider. (83) In general, the mother should avoid taking long-acting forms of medications and should watch her infant for any unusual signs and symptoms, such as sleepiness, irritability, or other known side effects of the prescribed medication. Acetaminophen and ibuprofen are well studied and safe. Some allergy medications can be sedating, so less sedating choices are better. Drugs contraindicated during breastfeeding include anti-cancer drugs, lithium, oral retinoids, iodine, amiodarone, and gold salts. (84) Online resources for guidance, such as LactMed, (85) InfantRisk Center (telephone 806/352-2519), (86) and the Postpartum Support International Warmline (telephone 800/944-4773) can help prevent giving incorrect advice about discontinuing a medication or discontinuing breastfeeding.

### Herbs or Supplements

Lactation specialists have recommended herbs such as fenugreek and blessed thistle, usually as a last resort when other nonpharmacologic measures do not result in an increase in milk volume. (87) However, milk expression is the

only successful, evidence-based method for increasing milk supply. During inquiry about maternal medication use, providers should question the mother about the use of herbal remedies or natural supplements. Appropriate counseling should be given to avoid the desire to take “magic bullets for milk supply.”

### Maternal Mental Health

Pregnancy-related depression and postpartum depression are associated with difficulty breastfeeding and shorter breastfeeding duration. Premature cessation of breastfeeding is also a risk factor for developing increased maternal anxiety. Women with high levels of anxiety and depression during pregnancy who stop breastfeeding early are at an additional multiplicative risk for postpartum anxiety and depression. (88) Best practice during a clinical visit is to screen the mother with a validated tool, such as the Edinburgh Postnatal Depression Scale, to ensure that maternal mental health is addressed adequately and objectively. (89)

### Alcohol, Tobacco, and Marijuana Use

Alcohol is transferred readily into human milk at levels that match the mother’s blood alcohol level. A safe rule, similar to the rule for safe drinking and driving a motor vehicle, is 1 drink consumed in 2 hours. Although alcohol test strips for breast milk are available for purchase, careful and limited consumption of alcohol is the best approach when breastfeeding. Tobacco smoking is not a contraindication to breastfeeding because the effects of smoking tobacco on the risk of sudden infant death syndrome and respiratory illness are almost negated if the infant has been breastfeeding. (90) However, nicotine can make an infant jittery, can interfere with let-down, and is associated with lower milk supply. Therefore, mothers who smoke should delay nursing as long as possible after smoking, and providers should provide motivational interviewing for smoking cessation assistance and/or suggest that a mother decrease the number of cigarettes she smokes or switch to cigarettes with lower nicotine content. Limited research is available on breastfeeding and marijuana use, including the amount of tetrahydrocannabinol (THC) in human milk, the length of time THC remains in the human milk after exposure, and the effects of THC on the infant. In a survey of more than 1,700 mothers in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) in Colorado, 6% reported using THC during pregnancy for symptoms of nausea, depression, and anxiety. (91) Because of concern for the developing brain in infants, mothers should abstain from using marijuana while breastfeeding. (92,93)

## Summary

- On the basis of strong evidence, the dose-responsive benefits of breastfeeding for both the infant and the mother are well established and should be the basis for recommending mothers to breastfeed exclusively. (25,26)
- On the basis of strong evidence, pain, sore nipples, and insufficient milk supply are the main reasons for early cessation of breastfeeding. (24,35)
- On the basis of strong evidence, hospital practices supportive of breastfeeding in the hours and days after birth make a difference in breastfeeding rates. However, currently, less than 20% of US births occur in Baby-Friendly Hospital Initiative–designated facilities.
- On the basis of consensus, in the first few weeks, stools commonly occur with every feeding or every other feeding. The “breastfed stool variant” occurs in about one-third of breastfed infants at about 4 to 6 weeks. One soft, voluminous stool occurs usually every 3 to 4 days, thought to be due to almost complete absorption of human milk.
- On the basis of consensus, developmental phases bring challenges for the breastfeeding dyad. Just when breastfeeding is getting established, infants go through a period of crying that may be misinterpreted as hunger and can lead to unnecessary supplementation. At about 4 months of age, most infants can become distracted during a feeding and are challenging to keep on the task of completing a full feeding. Nursing in a quiet, dark room with minimal distractions can help.
- On the basis of strong evidence and consensus, because milk supply is established in the first few weeks postpartum, access to trained internal staff or early referral for breastfeeding management and support is essential to ensure breastfeeding exclusivity. Hospital-based clinics or other maternal support groups provide additional support. (58,59)

- The breastfeeding touch points in Table 3 offer guidance for common issues that arise when supporting exclusive breastfeeding in the first 6 months postpartum and beyond.

To view teaching slides that accompany this article, visit <http://pedsinreview.aappublications.org/content/38/8/353.supplemental>.

### The Pediatrician’s Role in Encouraging Exclusive Breastfeeding

Maya Bunik, MD, MSPH

 Pediatrics in Review  
A PEDIATRIC PUBLICATION OF THE AMERICAN ACADEMY OF PEDIATRICS

American Academy of Pediatrics  
DEDICATED TO THE HEALTH OF ALL CHILDREN™

*References for this article are at <http://pedsinreview.aappublications.org/content/38/8/353>.*



## PIR Quiz

There are two ways to access the journal CME quizzes:

1. Individual CME quizzes are available via a handy blue CME link under the article title in the Table of Contents of any issue.
2. To access all CME articles, click "Journal CME" from Gateway's orange main menu or go directly to: <http://www.aapublications.org/content/journal-cme>.

1. A 22-year-old pregnant woman presents to the pediatrician's office at 37 weeks' gestation for a prenatal visit. She is currently interviewing the pediatric office practice in preparation for the birth of her first child. Although she intends to breastfeed, she is worried and has some concerns about sustaining breastfeeding as she works full-time and goes to school. She plans to take 6 weeks of maternity leave from work. At the prenatal visit, she asks the pediatrician for advice. Which of the following is the most appropriate office-based counseling that should be given to this patient?

- A. Advise her to breastfeed at night and supplement with formula during the day while she's away at work.
- B. Clearly communicate the importance of the exclusivity and duration of breastfeeding and assist her in working with her employer to facilitate this plan.
- C. Discourage her from starting breastfeeding if she is only going to do it for 6 weeks.
- D. Encourage her to discuss this topic with her obstetrician and the neonatologist in the nursery after delivery.
- E. Instruct her to supplement with formula right after delivery so that the baby gets used to the bottle, as she is unlikely to continue breastfeeding beyond 6 weeks.

2. You are performing an initial newborn examination on a 6-hour-old female newborn, following a full-term pregnancy and spontaneous vaginal delivery with no perinatal complications. You discuss with the parents the routine newborn anticipatory guidance messages. The counseling includes discussion of the benefits of human milk, particularly when it comes to boosting and supporting the immune system through the presence of several key proteins. Which of the following proteins has a lack of evidence to show a major role in the immunologic effects of breast milk?

- A. Casein.
- B. Complement.
- C. Cytokines.
- D. Lactoferrin.
- E. Lysozyme.

3. A gravida 1, para 1 woman gave birth to a 38-week-old female neonate and was discharged from the hospital 48 hours after delivery. She is exclusively breastfeeding. She brings her daughter to your clinic for her first visit. Today's weight is lower than the birth weight. The mother has several questions regarding the difference between breastfed versus formula-fed neonates. She is worried about the adequacy and variation in volume of her milk supply, the quality of her milk, and the baby's weight gain. Which of the following is the most appropriate response to address the concerns raised by this mother?

- A. Breastfed newborns regulate their milk intake to a greater degree than formula-fed newborns.
- B. Breast milk contains higher protein content than most formulas.
- C. In breastfed newborns who lose excessive weight in the first few weeks of life, supplementation is recommended to prevent dehydration and electrolyte imbalances.
- D. Newborns who are breastfed lose considerably less weight after birth than formula-fed newborns.
- E. Milk production is greatest in the morning, as prolactin and cortisol levels peak then.

**REQUIREMENTS:** Learners can take *Pediatrics in Review* quizzes and claim credit online only at: <http://pedsinreview.org>.

To successfully complete 2017 *Pediatrics in Review* articles for AMA PRA Category 1 Credit™, learners must demonstrate a minimum performance level of 60% or higher on this assessment, which measures achievement of the educational purpose and/or objectives of this activity. If you score less than 60% on the assessment, you will be given additional opportunities to answer questions until an overall 60% or greater score is achieved.

This journal-based CME activity is available through Dec. 31, 2019, however, credit will be recorded in the year in which the learner completes the quiz.



2017 *Pediatrics in Review* now is approved for a total of 30 Maintenance of Certification (MOC) Part 2 credits by the American Board of Pediatrics through the AAP MOC Portfolio Program. Complete the first 10 issues or a total of 30 quizzes of journal CME credits, achieve a 60% passing score on each, and start claiming MOC credits as early as October 2017.

4. The neonate in the previous question continues to receive routine follow-up at health supervision visits. She is brought to the clinic for a 6-month health supervision visit. The mother had successfully continued with exclusive breastfeeding. However, she reports that the baby has 2 teeth now, and a few weeks ago, the infant started biting at the nipples, and feedings are taking longer and are painful to her. Which of the following is the most appropriate explanation for the biting behavior in this patient?
- A. Biting allows the baby to control the pace of the transfer of the milk.
  - B. Biting at the beginning of the nursing session helps the baby complete the feeding in a shorter period of time.
  - C. Biting at the end of the nursing session may mean the baby is full and should end the feeding.
  - D. By 2 to 3 weeks of age, newborns may be distracted by the environment and take longer to feed, and biting help them focus on the feeding session.
  - E. This is a sign to completely discontinue breastfeeding and start formula.
5. A 3-month-old female infant who has been exclusively breastfed is brought to the pediatrician by her parents because of bloody stools. Other than slight fussiness, the baby has otherwise been feeding well, with no vomiting, fever, or change in activity level. Physical examination findings are unremarkable, with a benign abdominal examination finding and no evidence of anal fissure. The patient is given a presumptive diagnosis of allergic colitis. Which of the following is the most appropriate next step in the care of this patient?
- A. Discontinue breastfeeding and start regular formula feedings.
  - B. Eliminate cow milk products from the mother's diet and continue breastfeeding.
  - C. Start a course of oral prednisolone and continue breastfeeding.
  - D. Start the infant on lactose-free formula.
  - E. Start the infant on probiotic orally and continue breastfeeding.

## The Pediatrician's Role in Encouraging Exclusive Breastfeeding

Maya Bunik

*Pediatrics in Review* 2017;38;353

DOI: 10.1542/pir.2016-0109

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pedsinreview.aappublications.org/content/38/8/353>

### Supplementary Material

Supplementary material can be found at:  
<http://pedsinreview.aappublications.org/content/suppl/2017/07/31/38.8.353.DC1>  
<http://pedsinreview.aappublications.org/content/suppl/2017/08/21/38.8.353.DC2>

### References

This article cites 69 articles, 18 of which you can access for free at:  
<http://pedsinreview.aappublications.org/content/38/8/353.full#ref-list-1>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Medical Education**  
[http://classic.pedsinreview.aappublications.org/cgi/collection/medical\\_education\\_sub](http://classic.pedsinreview.aappublications.org/cgi/collection/medical_education_sub)  
**Journal CME**  
[http://classic.pedsinreview.aappublications.org/cgi/collection/journal\\_cme](http://classic.pedsinreview.aappublications.org/cgi/collection/journal_cme)  
**Nutrition**  
[http://classic.pedsinreview.aappublications.org/cgi/collection/nutrition\\_sub](http://classic.pedsinreview.aappublications.org/cgi/collection/nutrition_sub)  
**Breastfeeding**  
[http://classic.pedsinreview.aappublications.org/cgi/collection/breastfeeding\\_sub](http://classic.pedsinreview.aappublications.org/cgi/collection/breastfeeding_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<https://shop.aap.org/licensing-permissions/>

### Reprints

Information about ordering reprints can be found online:  
<http://classic.pedsinreview.aappublications.org/content/reprints>



## **The Pediatrician's Role in Encouraging Exclusive Breastfeeding**

Maya Bunik

*Pediatrics in Review* 2017;38;353

DOI: 10.1542/pir.2016-0109

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pedsinreview.aappublications.org/content/38/8/353>

Pediatrics in Review is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1979. Pediatrics in Review is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0191-9601.

# American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

